

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 16-3-98		3. REPORT TYPE AND DATES COVERED Technical Report
4. TITLE AND SUBTITLE The Dihydronaphthalene Elimination Reaction as a Route to Gallium-Nitrogen Compounds. Crystal and Molecular Structure of $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$			5. FUNDING NUMBERS Grant: N00014-96-1-0483 R&T Code: 4135002	
6. AUTHOR(S) O.T. Beachley, Jr., Matthew J. Noble, Melvyn Rowen Churchill and Charles H. Lake				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry State University of New York at Buffalo Natural Sciences Complex Buffalo, NY 14260-3000			8. PERFORMING ORGANIZATION REPORT NUMBER  Technical Report No. 51	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000			10. SPONSORING/MONITORING AGENCY REPORT NUMBER  n/a	
11. SUPPLEMENTARY NOTES  Accepted for publication - Organometallics				
12a. DISTRIBUTION/AVAILABILITY STATEMENT This document has been approved for public release and sale; its distribution is unlimited. Reproduction in whole or in part is permitted for any purpose of the United States government.			12b. DISTRIBUTION CODE  n/a	
13. ABSTRACT (Maximum 200 words)  The dihydronaphthalene derivative $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$ reacts at room temperature with $\text{NH}_3$ , $n\text{-PrNH}_2$ and $\text{PhNH}_2$ in THF solution to form high yields of $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNR}]_2$ ( $\text{R} = \text{H}$ , $n\text{-Pr}$ and $\text{Ph}$ ), $\text{C}_{10}\text{H}_{10}$ and $\text{NaCl}$ . In contrast, the elimination reactions between $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ with these same amines to form $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNR}]_2$ and $\text{PhCMe}_3$ require temperatures of $150^\circ\text{C}$ . The cyclopentadiene elimination reaction between $(\text{PhMe}_2\text{CCH}_2)_2\text{Ga}(\text{C}_5\text{H}_5)$ and aniline occurs at $-10^\circ\text{C}$ and is the fastest of these three. An X-ray structural study of $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$ identified it as the trans isomer.				
14. SUBJECT TERMS Organogallium compounds, elimination reactions, gallium amides			15. NUMBER OF PAGES 22	
			16. PRICE CODE n/a	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

19980615 032

OFFICE OF NAVAL RESEARCH

Contract N-00014-96-1-0483

R&T Code 4135002

TECHNICAL REPORT NO. 51

The Dihydronaphthalene Elimination Reaction as a Route to  
Gallium-Nitrogen Compounds. Crystal and Molecular Structure  
of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$

by

O. T. Beachley, Jr.\*, Matthew J. Noble,  
Melvyn Rowen Churchill\* and Charles H. Lake

Prepared for Publication

in

Organometallics

State University of New York at Buffalo  
Department of Chemistry  
Buffalo, New York 14260-3000

8 June 1998

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

\*This document has been approved for public release  
and sale; its distribution is unlimited

DTIC QUALITY INSPECTED 3

**The Dihydronaphthalene Elimination Reaction as a Route to  
Gallium-Nitrogen Compounds. Crystal and Molecular Structure  
of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$**

by

O. T. Beachley, Jr\*, Matthew J. Noble, Melvyn Rowen Churchill\* and Charles H. Lake

Department of Chemistry

State University of New York at Buffalo


Buffalo, NY 14260-3000

**Abstract**

The dihydronaphthalene derivative  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$  reacts at room temperature with  $\text{NH}_3$ ,  $n\text{-PrNH}_2$  and  $\text{PhNH}_2$  in THF solution to form high yields of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHR}]_2$  ( $\text{R} = \text{H}$ ,  $n\text{-Pr}$  and  $\text{Ph}$ ),  $\text{C}_{10}\text{H}_{10}$  and  $\text{NaCl}$ . In contrast, the elimination reactions between  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  with these same amines to form  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHR}]_2$  and  $\text{PhCMe}_3$  require temperatures of  $150^\circ\text{C}$ . The cyclopentadiene elimination reaction between  $(\text{PhMe}_2\text{CCH}_2)_2\text{Ga}(\text{C}_5\text{H}_5)$  and aniline occurs at  $-10^\circ\text{C}$  and is the fastest of these three. An X-ray structural study of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$  identified it as the trans isomer.


---

The hydrocarbon elimination reaction is of fundamental importance to group 13 - 15 chemistry.<sup>1</sup> In its simplest terms for nitrogen compounds, an amine adduct can be converted by an elimination reaction to an amide derivative, to an imide derivative and then





to a nitride. Researchers have attempted to define the factors which determine the temperatures necessary to initiate these reactions but without definitive success. Each succeeding reaction in this sequence typically requires higher temperatures. Since these reactions provide a direct connection between group 13 - 15 chemistry and the preparation of electronic materials such as GaN, InN, GaAs and InP, we have attempted to discover new hydrocarbon elimination reactions which occur at temperatures lower than those typically observed for the simple homoleptic organogallanes and group 15 bases ( $\text{ER}_{3-n}\text{H}_n$ , E = N, P, As; n = 1, 2, 3) because lower reaction temperatures minimize the possibility of decomposition reactions and, in turn, should lead to the formation of purer materials. This goal was realized with the discovery of the cyclopentadiene elimination reaction between  $\text{R}_2\text{Ga}(\text{C}_5\text{H}_5)$  (R =  $\text{Me}^2$ ,  $\text{Et}^3$  and  $\text{CH}_2\text{CMe}_3^4$ ) and amines and phosphines. These reactions typically occur below room temperature or 100 to 200 °C lower than the temperatures needed for reactions of  $\text{GaR}_3$  with the same bases. In this paper, elimination reactions of a diorganogallium derivative of dihydronaphthalene  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$  are described.

The yellow dihydronaphthalene derivative<sup>5, 6</sup>  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$




reacts at room temperature with  $\text{NH}_3$ ,  $n\text{-PrNH}_2$  and  $\text{PhNH}_2$  in THF solution to form  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHR}]_2$  ( $\text{R} = \text{H}$ ,  $n\text{-Pr}$  and  $\text{Ph}$ ) as colorless crystalline solids in high yields,  $\text{C}_{10}\text{H}_{10}$  and  $\text{NaCl}$  (eq 5). The time necessary for complete reaction was



indicated by the change in color of the solution from bright golden yellow to colorless at completion in all three cases. Aniline reacted fastest and required approximately 6 h for complete reaction whereas  $n$ -propylamine, the strongest base, was slowest and took 5 days. Ammonia required 4 days. The reaction of the neopentyl derivative  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$  with ammonia<sup>7</sup>, which was described previously, required only 18 h to go to completion. Thus, the neophyl derivative reacts significantly slower than the neopentyl derivative.

Since the dihydronaphthalene derivative  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$  decomposes at room temperature to form  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ,  $[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})]_n$  and  $\text{NaCl}$ , the question of whether  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHR}]_2$  ( $\text{R} = \text{H}$ ,  $n\text{-Pr}$  and  $\text{Ph}$ ) is formed by an elimination reaction between the amine and  $\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2]_2$  or from some other gallium compounds which could be in solution must be addressed. All experimental data are consistent with the conclusion that  $\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2]_2$  is the reactive compound. The simple elimination reactions of  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  with the amines were

studied independently and found to require temperatures in the order of 150 °C to initiate elimination (eq 6). No reaction was observed at room temperature. Second, if



$\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  had been formed from the decomposition of  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$ ,  $[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})]_n$ , a reddish brown compound<sup>5, 6</sup>, would also have been formed. However, the solution changed from lemon yellow to bright golden yellow to colorless and never was reddish brown. These observations suggest that  $[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})]_n$  was not present. Furthermore, the closely related low oxidation state gallium compound  $[\text{Ga}(\text{CH}_2\text{CMe}_3)]_n$  has been reacted with ammonia<sup>7</sup> but temperatures around 450 °C were required to initiate a reaction. The products were GaN, H<sub>2</sub> and CMe<sub>4</sub>. Third, the initial lemon yellow color is due to  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$ . When the amine was added, a colorless precipitate (assumed to be NaCl) was formed and the color of the solution changed to bright golden yellow. These observations are consistent with the formation of an amine adduct,  $\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2]_2 \cdot 2\text{amine}$ , the species which eventually is transformed into  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHR}]_2$  and  $\text{C}_{10}\text{H}_{10}$ .

The dihydronaphthalene elimination reaction between  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$  and aniline was compared directly with the cyclopentadiene elimination reaction between  $(\text{PhMe}_2\text{CCH}_2)_2\text{Ga}(\text{C}_5\text{H}_5)$  and aniline. The latter was prepared *in situ* by a stoichiometric ligand redistribution reaction between  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  and  $\text{Ga}(\text{C}_5\text{H}_5)_3$ .<sup>4</sup> After a stoichiometric amount of aniline was added to the solution of  $(\text{PhMe}_2\text{CCH}_2)_2\text{Ga}(\text{C}_5\text{H}_5)$ , the reaction mixture was maintained at -10 °C and after 24 h  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$ , a slightly soluble solid, was isolated in higher than 70 % yield. Thus, the unsymmetrically substituted neophylgallium compounds  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaC}_5\text{H}_5$  and  $[(\text{PhMe}_2\text{CCH}_2)_2\text{Ga}]_2(\text{C}_{10}\text{H}_8)$  both undergo elimination

reactions with amines at room temperature (or below) to form  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHR}]_n$ , but the cyclopentadiene elimination reaction is faster.

Three gallium nitrogen compounds  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHR}]_2$  ( $\text{R} = \text{H}$ ,  $n\text{-Pr}$ <sup>8</sup> and  $\text{Ph}$ ) were prepared and characterized in this study but only  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$  was characterized by an X-ray structural study. The products formed at room temperature have sharper and slightly higher melting points and are probably more pure than those formed in sealed tubes at 140 - 150 °C. Cryoscopic molecular weight data demonstrated the presence of dimers. <sup>1</sup>H NMR spectral data show the trans isomer being the predominant species in benzene solution. The cis isomer was observed in 7 % abundance but only in those benzene solutions of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$  which had been prepared from the dihydronaphthalene derivative at room temperature. A benzene solution of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$  which had been prepared in a sealed tube at 140 - 150 °C consisted of 100 % trans isomer. When the <sup>1</sup>H NMR spectrum of a  $\text{CH}_2\text{Cl}_2$  solution of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{Ga-NHPh}]_2$  which had been prepared by the dihydronaphthalene elimination reaction was recorded, 16 % cis isomer was observed. Benzene solutions of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNH}(n\text{-Pr})]_2$  prepared by either route contained only the 100 % trans isomer.

An X-ray structural study of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$  showed the crystal to be composed solely of the trans isomer. The crystallographic asymmetric unit consists of one-half of each of two crystallographically independent molecules, each of which lies about a center of symmetry. Molecule 1 lies about 0, 1/2, 0 (Wyckoff notation c), while molecule 2 lies about 1/2, 1/2, 1/2 (Wyckoff notation h).<sup>9</sup> All molecules are separated by normal van der Waals' distances and there are no anomalously short intermolecular contacts (Figure 1). The labeling scheme used for "Molecule 1" is shown in Figure 2. Note that the two crystallographically independent molecules are chemically equivalent and vary only in minor differences of orientation about Ga-C or Ga-N linkages. Bond distances and angles for both molecules are presented in Tables 1 and 2, respectively.

The presence of centers of symmetry necessitates that each  $\text{Ga}_2\text{N}_2$  molecular core be strictly planar. Each four-membered ring is slightly distorted from its possible  $D_{2h}$  symmetry. Gallium-nitrogen bond lengths range from 2.037(3) Å to 2.082(4) Å. The cross-ring distances associated with the  $\text{Ga}_2\text{N}_2$  molecular core are  $\text{Ga}(1)\cdots\text{Ga}(1A) = 3.008$  Å and  $\text{N}(3)\cdots\text{N}(3A) = 2.815$  Å (3.018 Å and 2.780 Å, respectively, for the second molecule). These data may be compared with the corresponding data for the closely related compound  $[(\text{PhMe}_2\text{CCH}_2)_2\text{Ga-NH}(\text{n-Pr})]_2$ .<sup>8</sup> The Ga-N distances were slightly shorter at 2.013(2) and 2.029(2) Å and the  $\text{Ga}_2\text{N}_2$  molecular core was slightly more compact with the two Ga atoms separated by 2.938(1) Å and the two nitrogen atoms by 2.776(3) Å. The internal ring angle about Ga was 86.7(1)° whereas about N it was 93.3(1)°.

Two neophyl ligands are bound to each gallium(III) center with  $\text{Ga}(1)\text{-C}(00) = 1.967(6)$  Å and  $\text{Ga}(1)\text{-C}(10) = 1.982(5)$  Å and an interligand angle of  $\text{C}(00)\text{-Ga}(2)\text{-C}(10) = 123.5(2)^\circ$  ( $\text{Ga}(2)\text{-C}(30) = 1.988(5)$  Å,  $\text{Ga}(2)\text{-C}(40) = 1.982(5)$  Å and  $\text{C}(30)\text{-Ga}(2)\text{-C}(40) = 122.2(2)^\circ$  the second molecule). The Ga-C distances for  $[(\text{PhMe}_2\text{CCH}_2)_2\text{Ga-NH}(\text{n-Pr})]_2$ <sup>8</sup> were 1.992(2) and 1.994(3) Å. The twist associated with the neophyl ligands for  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$  in relation to the molecular core is 89.3° as defined by the intercept of the planes  $\text{C}(00)\text{-Ga}(1)\text{-C}(10)$  and  $\text{N}(3)\text{-Ga}(1)\text{-N}(3A)$  ( $\text{C}(30)\text{-Ga}(2)\text{-C}(40)/\text{N}(4)\text{-Ga}(2)\text{-N}(4A) = 88.1^\circ$  for the second molecule). The  $\alpha$ -carbon atoms of the neophyl ligands possess a distorted tetrahedral environment with expanded  $\text{Ga-C}(\alpha)\text{-C}(\beta)$  angles of 116.6(3) to 119.9(4)°.

Primary phenyl amide ligands bridge the gallium(III) metal centers. The nitrogen atoms of these ligands possess a distorted tetrahedral environment (Table 2, Section b). The hydrogen atoms of the amine moieties were located and refined successfully yielding N-H distances of 0.89(6) Å and 0.81(5) Å, respectively, for H(3) and H(4). The N-H distance in  $[(\text{PhMe}_2\text{CCH}_2)_2\text{Ga-NH}(\text{n-Pr})]_2$  was 0.82(3) Å.<sup>8</sup> The phenyl ring is bonded to the nitrogen atom with  $\text{N}(3)\text{-C}(21) = 1.437(8)$  Å ( $\text{N}(4)\text{-C}(51) = 1.436(7)$  Å. The phenyl rings intercept the plane of the molecular core at 85.8° and 81.6° for molecules 1 and 2,



respectively (intercept of the planes Ga(1)-N(3)-Ga(1A)/ C(21)-C(22)-C(23) and Ga(2)-N(4)-Ga(2A)/ C(51)-C(52)-C(53)). The phenyl rings are close to parallel and are perpendicular to the appropriate Ga...Ga axis, an orientation resulting from the arrangement of the neophyl ligands about the molecular core.

### Experimental Section

All compounds described in this investigation were very sensitive to oxygen and moisture and were manipulated under a purified argon atmosphere in a Vacuum Atmospheres drybox equipped with a Dry Train or by using standard vacuum line techniques. The reagents  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  and  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$  were prepared by using literature procedures.<sup>8</sup> All solvents were purified before use. Elemental analyses were performed by E+R Microanalytical Laboratories, Corona, NY. The  $^1\text{H}$  NMR spectra were recorded at 400 MHz by using a Varian VXR-400 spectrometer with samples in flame-sealed NMR tubes. Chemical shifts are reported in  $\delta$  (ppm) and are referenced to tetramethylsilane (TMS) at  $\delta = 0.00$  ppm and benzene at  $\delta = 7.15$  ppm. Infrared spectra of Nujol mulls between CsI plates were recorded by means of a Perkin Elmer 683 spectrometer. Melting points were observed with a Mel-Temp by using flame-sealed capillaries and are uncorrected. Molecular weights were measured cryoscopically in benzene by using an instrument similar to that described by Shriver and Drezdson.<sup>10</sup>

**Reaction of  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$  with Anhydrous Amines.** In a typical experiment the yellow gallium(III) dihydronaphthalene derivative<sup>5, 6</sup>  $\text{Na}_2\{\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2\}$  was prepared in a Solv-Seal reaction flask at  $-78^\circ\text{C}$  from sodium metal, naphthalene,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$  and 40 - 50 mL of THF. Then, a slight excess of the dry amine was added to the flask by vacuum distillation and the solution was warmed from  $-196^\circ\text{C}$  to  $-78^\circ\text{C}$ . A large amount of colorless precipitate (NaCl) formed as the solution became bright yellow. Finally, the reaction mixture was warmed with stirring to ambient temperature over 12 h. When reaction with the amine was complete, the mixture was colorless. THF and the other readily volatile materials were

removed by vacuum distillation at 20 °C. After the bulk of the readily volatile material had been removed, the material remaining in the flask was subjected to dynamic vacuum for 24 h. The slightly volatile materials were collected in a small preweighed trap and the components were identified by  $^1\text{H}$  NMR spectroscopy as  $\text{C}_{10}\text{H}_{10}$ ,  $\text{C}_{10}\text{H}_8$ , and THF. The nonvolatile material was extracted with pentane in order to separate NaCl from the soluble gallium-nitrogen product. The latter was purified and then fully characterized. The following paragraph gives quantities of reactants, the time required at room temperature for complete reaction, quantities and percent yields of products and the characterization data for each colorless new compound.

**$[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNH}_2]_2$ .** Reagents: 0.0926 g Na (4.03 mmol), 0.517 g  $\text{C}_{10}\text{H}_8$  (4.03 mmol); 1.49 g  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$  (4.00 mmol), 4.22 mmol  $\text{NH}_3$ . Time at 20 °C for complete reaction: 4 d. Products: 0.231 g NaCl (3.95 mmol, 98.9 % yield based on Na, see Eq 1), 1.30 g  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaNH}_2$  (3.69 mmol, 92.2 % yield based on  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$  (Eq 1), product purified by recrystallization from pentane at -30 °C). Characterization data: Mp: 99.1 - 101.1 °C.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ,  $\delta$ ): 7.16 (d, 4 H, *o*-Ar,  $^3J_{\text{CCH}} = 7.59$  Hz), 7.04 (t, 4 H, *m*-Ar,  $^3J_{\text{CCH}} = 6.26$  Hz), 6.90 (t, 2 H, *p*-Ar,  $^3J_{\text{CCH}} = 6.33$  Hz), 1.20 (s, 12 H, -CMe<sub>2</sub>), 0.53 (s, 4 H, -CH<sub>2</sub>-), -0.94 (s, 1 H, -NH<sub>2</sub>). IR (Nujol,  $\text{cm}^{-1}$ ): 3378 (vs), 3310 (s), 3080 (s), 3075 (s), 3019 (s), 1951 (sh, m), 1943 (m), 1882 (m), 1867 (m), 1800 (m), 1744 (m), 1670 (m), 1597 (s), 1587 (m), 1275 (s), 1178 (s), 1155 (m), 1130 (m), 1098 (m), 1069 (s), 1061 (sh,s), 1027 (s), 998 (m), 960 (m), 942 (m), 925 (m), 901 (m), 848 (m), 848 (m), 838 (s), 821 (vs), 775 (m), 762 (vs), 715 (vs), 692 (vs), 635 (s), 615 (sh, m), 605 (sh, m), 585 (m), 554 (s), 508 (w), 445 (s), 269 (w). Anal. for  $\text{C}_{20}\text{H}_{28}\text{GaN}$ . Calcd: C, 68.21; H, 8.01. Found: C, 68.14; H, 7.98. Solubility: soluble in THF, pentane, and benzene.

**$[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNH}(\text{n-Pr})]_2$ .** Reagents: 0.117 g Na (5.09 mmol), 0.653 g  $\text{C}_{10}\text{H}_8$  (5.09 mmol); 1.86 g  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$  (5.00 mmol), 0.301 g  $\text{n-PrNH}_2$  (5.09 mmol). Time at 20 °C for complete reaction: 5 days. Products: 0.264 g NaCl (4.52 mmol,

90.3 % yield based on Na and Eq 1), 1.67 g  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{n-Pr})$  (4.23 mmol, 84.6 % yield based on  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$ , after product purified by recrystallization from pentane at  $-30^\circ\text{C}$ ). Characterization data: Mp:  $127.8 - 128.5^\circ\text{C}$  (lit<sup>8</sup>  $129.3 - 131.0^\circ\text{C}$ ).  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ,  $\delta$ ): 7.35 (d, 4 H, *o*-Ar,  $^3J_{\text{CCH}} = 8.10$  Hz), 7.16 (t, 4 H, *m*-Ar,  $^3J_{\text{CCH}} = 7.26$  Hz), 7.00 (t, 2 H, *p*-Ar,  $^3J_{\text{CCH}} = 7.14$  Hz), 2.06 (q, 2 H,  $-\text{CH}_2-$  (Pr),  $^3J_{\text{CCH}} = 7.68$  Hz), 1.33 (s, 12 H,  $-\text{CMe}_2$  (neophyl)), 0.87 (s, 4 H,  $-\text{CH}_2-$  (neophyl)), 0.82 (m, 2 H,  $-\text{NCH}_2-$  (Pr)), 0.61 (t, 3 H,  $-\text{CH}_3$  (Pr),  $^3J_{\text{CCH}} = 7.14$  Hz), - 0.04 (t, 1 H,  $-\text{NH}$ ,  $^3J_{\text{CNH}} = 7.50$  Hz). The  $^1\text{H}$  NMR spectrum indicated 100% trans isomer. Solubility: soluble in THF, pentane, and benzene.

**$[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$ :** Reagents: 0.115 g Na (5.01 mmol), 0.643 g  $\text{C}_{10}\text{H}_8$  (5.01 mmol); 1.86 g  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$  (5.00 mmol), 0.480 g  $\text{PhNH}_2$  ( 5.15 mmol). Time at  $20^\circ\text{C}$  for complete reaction: 6 h. Products: 0.272 g NaCl (4.65 mmol, 93.1 % yield based on Na and Eq 1), 1.75 g  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}$  (4.08 mmol, 81.6 % yield based on  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$  after product purified by recrystallization from toluene at  $-78^\circ\text{C}$ ). Characterization data: Mp:  $175.4 - 177.5^\circ\text{C}$ .  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ,  $\delta$ ) 7.30 (d, 8 H, *o*-Ar (neophyl),  $^3J_{\text{CCH}} = 7.60$  Hz), 7.20 (t, 4 H, *m*-Ar (N-Ph),  $^3J_{\text{CCH}} = 6.80$  Hz), 7.17 (t, 8 H, *m*-Ar (neophyl),  $^3J_{\text{CCH}} = 7.40$  Hz), 7.03 (t, 4 H, *p*-Ar (neophyl),  $^3J_{\text{CCH}} = 7.20$  Hz), 6.85 (t, 2 H, *p*-Ar (N-Ph),  $^3J_{\text{CCH}} = 7.40$  Hz), 6.78 (d, 4 H, *o*-Ar (N-Ph),  $^3J_{\text{CCH}} = 8.80$  Hz), 3.15 (s-br, 2.1 H,  $-\text{NH}$  (trans + cis)), 1.07 (d, 4.0 H,  $-\text{CH}_2-$  (trans),  $^2J_{\text{CH}} = 13.6$  Hz), 1.03 (s, 12.0 H,  $-\text{CMe}_2$  (trans)), 1.01 (s, 12.9 H,  $-\text{CMe}_2$  (trans + cis)), 0.77 (d, 3.8 H,  $-\text{CH}_2-$  (trans),  $^2J_{\text{CH}} = 13.6$  Hz). The  $^1\text{H}$  NMR spectrum for a  $\text{C}_6\text{D}_6$  solution indicated an isomer distribution of 7% cis and 93% trans ( $\pm 5\%$ ).  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ ,  $\delta$ ): 7.34, 7.33, 7.30, 7.28, 7.25, 7.15 - (m, 24 H, *o*-Ar (neophyl), *m*-Ar (neophyl), *p*-Ar (neophyl), *m*-Ar (N-Ph)), 6.99 (t, 2 H, *p*-Ar (N-Ph),  $^3J_{\text{CCH}} = 6.00$  Hz), 6.55 (d, 4 H, *o*-Ar (N-Ph),  $^3J_{\text{CCH}} = 7.30$  Hz), 2.94 (s-br, 2.5 H,  $-\text{NH}$  (trans + cis)), 1.22 (s, 2.30 H,  $-\text{CMe}_2$  (cis)), 1.02 (s, 0.70 H,  $-\text{CH}_2-$  (cis)), 0.96 (s, 25.9 H,  $-\text{CMe}_2$  (trans + cis)), 0.87 (d, 4.2 H,  $-\text{CH}_2-$  (trans),  $^2J_{\text{CH}} = 13.7$  Hz), 0.82 (s, 0.70 H,  $-\text{CH}_2-$  (cis)), 0.58 (d, 4.2 H,  $-\text{CH}_2-$  (trans),

$^2J_{\text{CH}} = 14.3$  Hz). The  $^1\text{H}$  NMR spectrum for the  $\text{CH}_2\text{Cl}_2$  solution indicated an isomer distribution of 16% cis and 84% trans ( $\pm 5\%$ ). IR (Nujol mull,  $\text{cm}^{-1}$ ): 3275 (s), 3082 (s), 3055 (s), 3015 (s), 2721 (m), 1949 (m), 1871 (m), 1809 (m), 1750 (m), 1595 (s), 1585 (sh, s), 1575 (m), 1489 (vs), 1279 (m), 1231 (s), 1210 (vs), 1175 (s), 1155 (m), 1139 (m), 1100 (m), 1070 (s), 1025 (s), 998 (m), 963 (m), 905 (m), 892 (s), 868 (sh, m), 862 (s), 852 (sh, m), 839 (m), 811 (m), 761 (vs), 750 (s), 725 (s), 712 (m), 699 (sh, vs), 692 (vs), 681 (sh, s), 665 (sh, m), 626 (m), 602 (w), 569 (m), 555 (m), 535 (w), 512 (w), 430 (sh, w), 418 (m), 408 (m), 250 (w), 235 (m). Anal. for  $\text{C}_{26}\text{H}_{32}\text{GaN}$ . Calcd: C, 72.92; H, 7.53. Found: C, 72.82; H, 7.59. Solubility: soluble in THF; slightly soluble in pentane, benzene, toluene and *m*-xylene. Compound had insufficient solubility in benzene for a cryoscopic molecular weight study.

**Reactions of  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  with  $\text{NH}_3$  and with  $\text{PhNH}_2$ .** In a typical experiment a break-seal reaction tube was charged with  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  and the amine, evacuated, sealed and then heated for 4 days at 140-150  $^\circ\text{C}$ . After no noncondensable gas was observed to have been formed during the reaction by using the vacuum line, the more volatile product  $\text{C}_6\text{H}_5\text{CMe}_3$  was transferred to an NMR tube by vacuum distillation at ambient temperature and identified as *t*-butylbenzene by  $^1\text{H}$  NMR spectroscopy. The remaining colorless solid was washed out of the reaction tube with  $\sim 50$  mL of pentane through a medium glass frit, purified by recrystallization from pentane and identified.

**$[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNH}_2]_2$ .** Reagents: 2.31 g  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  (4.93 mmol);  $\text{NH}_3$  (5.01 mmol). Products: 0.317 g  $\text{C}_6\text{H}_5\text{CMe}_3$  (4.40 mmol, 93.2% yield based on  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ), 1.64 g  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaNH}_2$  (4.64 mmol, 94.2% based on  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ). Characterization data: Mp: 94.3 - 97.3  $^\circ\text{C}$ . Anal. calcd for  $\text{C}_{20}\text{H}_{28}\text{GaN}$ : C, 68.21; H, 8.01. Found: C, 68.48; H, 7.96. Cryoscopic molecular weight, benzene solution, formula weight 352.17 (obsd. molality, obsd. mol. wt., association): 0.0909, 686, 1.95; 0.0721, 695, 1.98; 0.0550, 708, 2.01. Spectroscopic data for this sample were identical within normal experimental error with the data for the compound prepared by the

reaction of  $\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2]_2$  with  $\text{NH}_3$ . Solubility: soluble in pentane and benzene.

**$[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$ :** Reagents: 2.20 g  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  (4.69 mmol); 0.503 g  $\text{PhNH}_2$  (5.40 mmol). Products: 1.36 g  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}$  (3.18 mmol, 67.8% yield based on  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ). Characterization data: Mp: 174.1 - 177.0 °C. Spectroscopic data for this sample were identical within normal experimental error with the data for the compound prepared from the reaction of  $\text{C}_{10}\text{H}_8[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2]_2$  with  $\text{PhNH}_2$ . Only the trans isomer was observed by  $^1\text{H}$  NMR spectroscopy.

**Synthesis of  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}$  by Reacting  $\text{Ga}(\text{C}_5\text{H}_5)_3$ ,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  and  $\text{PhNH}_2$ .** A reaction flask was charged with  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  (1.46 g, 3.11 mmol),  $\text{Ga}(\text{C}_5\text{H}_5)_3$  (0.411 g, 1.55 mmol) and 20 - 25 mL of pentane. After the solution had been stirred for 1 h at 20 °C, the flask was cooled to -10 °C and a solution of  $\text{PhNH}_2$  (0.434 g, 4.66 mmol) in 20 mL of pentane was added slowly. This solution was stirred for an additional 24 h and a large amount of colorless precipitate, the slightly soluble product, formed. After the pentane was removed by vacuum distillation, the residue was washed numerous times with fresh pentane at low temperature (-40 to -78 °C) to remove impurities. The resulting colorless solid was identified as pure  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}$  (1.41 g, 3.30 mmol, 70.8% yield based on  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ). Crystallographic quality crystals were obtained by slowly cooling to ambient temperature a heated, saturated solution of the compound in *m*-xylene. The crystals were mounted directly from the *m*-xylene solution.

**$[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$ :** Mp: 173.6 - 176.0 °C.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ,  $\delta$ ) (400 MHz): 7.28 (d, 8.0 H, *o*-Ar (neophyl),  $^3J_{\text{CCH}} = 8.0$  Hz), 7.19 (t, 4.0 H, *m*-Ar (N-Ph),  $^3J_{\text{CCH}} = 7.80$  Hz), 7.17 (t, 8.0 H, *m*-Ar (neophyl),  $^3J_{\text{CCH}} = 7.80$  Hz), 7.02 (t, 4.0 H, *p*-Ar (neophyl),  $^3J_{\text{CCH}} = 7.40$  Hz), 6.84 (t, 2.0 H, *p*-Ar (N-Ph),  $^3J_{\text{CCH}} = 7.40$  Hz), 6.76 (d, 4.0 H, *o*-Ar (N-Ph),  $^3J_{\text{CCH}} = 8.00$  Hz), 3.14 (s-br, 2.0 H, -NH (trans)), 1.06 (d, 3.9 H, -

CH<sub>2</sub>- (trans),  $^2J_{\text{CH}} = 14.0$  Hz), 1.02 (s, 11.8 H, -CMe<sub>2</sub> (trans)), 1.00 (s, 11.8 H, -CMe<sub>2</sub> (trans)), 0.76 (d, 3.9 H, -CH<sub>2</sub>- (trans),  $^2J_{\text{CH}} = 13.6$  Hz). Only the trans isomer was observed by <sup>1</sup>H NMR spectroscopy. Anal. for C<sub>26</sub>H<sub>32</sub>GaN. Calcd.: C, 72.92; H, 7.53. Found: C, 72.93; H, 7.44. Solubility: slightly soluble in pentane, benzene, and *m*-xylene.

**Collection of X-Ray Diffraction Data for [(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaNHPh]<sub>2</sub>.**

A well-defined crystal of approximate orthogonal dimensions 0.30 × 0.25 × 0.15 mm was sealed into a thin-walled glass capillary under an argon atmosphere inside the drybox maintained under anaerobic and moisture-free conditions. The crystal was inspected under a binocular microscope to ensure that it was indeed a single crystal and then was centered on a Siemens R3m/V automated four-circle diffractometer. The unit cell parameters were determined as described previously.<sup>11</sup> Intensity data (Mo Kα radiation,  $\lambda = 0.710730 \text{ \AA}$ )<sup>12</sup> were collected at ambient temperature (25 ± 1 °C) with graphite-monochromatized radiation. The observed Laue symmetry (C<sub>i</sub> or  $\bar{1}$  only) and the non-occurrence of systematic absences indicate that the crystal belongs to the triclinic system. The centrosymmetric space group  $P\bar{1}$  (No. 2) was selected on the basis of intensity statistics and frequency of occurrence<sup>13</sup> and was confirmed by the successful solution of the structure. A total of 6373 reflections were collected and merged to yield 6004 unique reflections of which 3711 (61.8%) had  $|F_o| > 6.0\sigma(F_o)$ . Details appear in Table 3.

**Determination of Crystal Structure of [(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaNHPh]<sub>2</sub>.** All crystallographic calculations were carried out on a VAX station computer with the use of the Siemens SHELXTL PLUS (Release 4.11 (VMS)) program package.<sup>14, 15</sup> The analytical expressions of the scattering factors for neutral atoms<sup>16a</sup> were corrected for the  $\Delta f'$  and  $i\Delta f''$  components of anomalous dispersion.<sup>16b</sup> The structure was solved via a combination of Patterson maps, difference-Fourier syntheses and least-squares refinement. Positional and anisotropic thermal parameters for the unique N-bonded hydrogen atom of each independent phenylamido group were also refined; all remaining hydrogen atoms were placed in calculated positions, based upon  $d(\text{C-H}) = 0.96 \text{ \AA}$ <sup>17</sup> and the appropriate idealized

trigonal or staggered tetrahedral geometry. Refinement of the ordered model converged with  $R = 3.43\%$  and  $wR = 3.62\%$  for those 3711 data with  $|F_o| > 6.0\sigma(|F_o|)$ , and  $R = 6.94\%$  for all data.

**Acknowledgment.** This work was supported in part by the Office of Naval Research (O.T.B.). The purchase of the X-ray diffractometer was made possible by Grant 89-13733 from the Chemical Instrumentation Program of the National Science Foundation.

**Supporting Information Available.** One Figure (showing labeling of atoms in "Molecule 2"); complete tables of positional parameters, interatomic distances and angles, anisotropic thermal parameters, and calculated positions for hydrogen atoms for the X-ray study (11 pp). Ordering information is given on any current masthead page.

## References

1. (a) Wilkinson, G.; Stone, F. G. A., Abel, E.W., Eds. *Comprehensive Organometallic Chemistry*; Pergamon Press: Oxford, 1982; Vol.1, p 588 and references therein.  
 (b) Coates, G. E.; Green, M. L. H.; Wade, K., Eds. *Organometallic Compounds*; Methuen: London, 1967; Vol. 1, p 177 and references therein. (c) Cowley, A. H.; Jones, R. H. *Angew. Chem., Int. Ed. Engl.* **1989**, 28, 1208. (d) Schauer, S. J.; Lake, C. H.; Watkins, C. L.; Krannich, L. K.; Powell, D. H. J. *Organomet. Chem.* **1997**, 549, 31 and references therein.
2. Beachley, O. T., Jr.; Royster, T. L., Jr.; Arhar, J. R.; Rheingold, A. L. *Organometallics* **1993**, 12, 1976.
3. Beachley, O. T., Jr.; Rosenblum, D. B.; Churchill, M. R.; Lake, C. H.; Toomey, L. *M. Organometallics* **1996**, 15, 3653.
4. Beachley, O. T., Jr.; Maloney, J. P.; Rogers, R. D. *Organometallics* **1997**, 16, 3267.
5. Noble, M. J.; State University of New York at Buffalo; Ph. D. Thesis; 1994.
6. Beachley, O. T., Jr.; Pazik, J. C.; Noble, M. J. *Organometallics* **1994**, 13, 2885.
7. Beachley, O. T., Jr.; Pazik, J. C.; Noble, M. J. *Organometallics* **1998**, 17, 0000.
8. Beachley, O. T., Jr.; Noble, M. J.; Churchill, M. R.; Lake, C. H. *Organometallics* **1992**, 11, 1051.
9. (a) International Tables for X-Ray Crystallography, Kynoch Press, Birmingham, England, **1965**, Vol. 1 (2nd Ed.), p 75. (b) A reviewer suggested the possibility of "a monoclinic C-centered cell (11.054, 20.365, 20.502 Å, beta [sic] = 96.5 °". If one doubles the b-axis in our present triclinic unit cell, there results a C-centered cell with  $a' = 11.0541(19)$ ,  $b' = 20.3671(44)$ ,  $c' = 20.5098(36)$  Å,  $\alpha' = 90.321(15)$ ,  $\beta' = 96.486(16)$ ,  $\gamma' = 91.587(16)$  °. Both  $\alpha'$  and  $\gamma'$  are significantly different from 90 °, and there is no evidence for 2/m symmetry of diffraction. Furthermore, the completed structure (viz. Figure 1) shows no general crystallographic relationship between "Molecule 1" and "Molecule 2". The attribution of our crystal to the triclinic crystal class



is thus secure.

10. Shriver, D. F.; Drezdson, M. A. *The Manipulation of Air Sensitive Compounds*; Wiley: New York, 1986; p 38.
11. Churchill, M. R.; Lashewycz, R. A.; Rotella, F. J. *Inorg. Chem.* **1977**, *16*, 265.
12. Bearden, J. A. *Rev. Mod. Phys.* **1967**, *39*, 78.
13. Nowacki, W.; Matsumato, T.; Edenharter, A. *Acta Crystallogr.* **1967**, *22*, 935.
14. Sheldrick, G. M. SHELXTL PLUS, Release 4.11 (VMS), Siemens Analytical Instrument Corp., Madison, WI, **1989**.
15. Sheldrick, G. M. SHELXTL PLUS, An Integrated System for Solving, Refining and Displaying Crystal Structures from Diffraction Data; For Nicolet R3m/V; University of Göttingen, Germany, **1987**.
16. International Tables for X-Ray Crystallography, Kynoch Press, Birmingham, England, **1974**, Vol. 4 (a) pp 99-101; (b) pp 149-150.
17. Churchill, M. R. *Inorg. Chem.* **1973**, *12*, 1213.

Table 1

Selected Bond Distances (Å) for  $[(\text{PhMe}_2\text{CCH}_2)\text{GaNHPh}]_2$ .

## MOLECULE 1

## MOLECULE 2

## (a) Distances Associated with the Gallium(III) Atoms

Ga(1)–N(3)	2.082(4)	Ga(2)–N(4)	2.066(4)
Ga(1)–C(00)	1.977(6)	Ga(2)–C(30)	1.988(5)
Ga(1)–C(10)	1.982(5)	Ga(2)–C(40)	1.982(5)
Ga(1)···Ga(1A)	3.008(1)	Ga(2)···Ga(2A)	3.018(1)
Ga(1)–N(3A)	2.037(3)	Ga(2)–N(4A)	2.037(5)

## (b) Bond Lengths within the Bridging Amide Ligands

N(3)–H(3)	0.89(6)	N(4)–H(4)	0.81(5)
N(3)–C(21)	1.437(8)	N(4)–C(51)	1.436(7)
N(3)–Ga(1A)	2.037(3)	N(4)–Ga(2A)	2.037(5)
C(21)–C(22)	1.375(9)	C(51)–C(52)	1.365(7)
C(21)–C(26)	1.377(6)	C(51)–C(56)	1.392(9)
C(22)–C(23)	1.389(10)	C(52)–C(53)	1.397(10)
C(23)–C(24)	1.392(8)	C(53)–C(54)	1.370(13)
C(24)–C(25)	1.358(13)	C(54)–C(55)	1.365(9)
C(25)–C(26)	1.368(10)	C(55)–C(56)	1.375(9)

Table 2

Selected Bond Angles (deg) for  $[(\text{PhMe}_2\text{CCH}_2)\text{GaNHPh}]_2$ .

## MOLECULE 1

## MOLECULE 2

## (a) Angles around the gallium(III) centers

N(3)–Ga(1)–C(00)	108.5(2)	N(4)–Ga(2)–C(30)	106.1(2)
N(3)–Ga(1)–C(10)	107.8(2)	N(4)–Ga(2)–C(40)	109.9(2)
C(00)–Ga(1)–C(10)	123.5(2)	C(30)–Ga(2)–C(40)	122.2(2)
N(3)–Ga(1)–N(3A)	86.2(1)	N(4)–Ga(2)–N(4A)	85.3(2)
C(00)–Ga(1)–N(3A)	111.5(2)	C(30)–Ga(2)–N(4A)	113.6(2)
C(10)–Ga(1)–N(3A)	112.6(2)	C(40)–Ga(2)–N(4A)	113.0(2)

## (b) Angles around the nitrogen atoms

Ga(1)–N(3)–Ga(1A)	93.8(1)	Ga(2)–N(4)–Ga(2A)	94.7(2)
Ga(1)–N(3)–C(21)	117.6(3)	Ga(2)–N(4)–C(51)	118.7(3)
Ga(1A)–N(3)–C(21)	125.2(3)	Ga(2A)–N(4)–C(51)	123.8(3)
Ga(1)–N(3)–H(3)	96(3)	Ga(2)–N(4)–H(4)	102(3)
Ga(1A)–N(3)–H(3)	113(4)	Ga(2A)–N(4)–H(4)	107(4)
C(21)–N(3)–H(3)	107(4)	C(51)–N(4)–H(4)	108(4)

(c) Angles about C( $\alpha$ ) of the neophyl groups

Ga(1)–C(00)–C(01)	119.9(4)	Ga(2)–C(30)–C(31)	116.6(3)
Ga(1)–C(10)–C(11)	118.3(4)	Ga(2)–C(40)–C(41)	119.3(4)

Table 3.

Data for the X-Ray Crystallographic Studies of  $[(\text{PhMe}_2\text{CCH}_2)\text{GaNHPh}]_2$ 

mol formula	$\text{C}_{52}\text{H}_{64}\text{Ga}_2\text{N}_2$
cryst syst	triclinic
space Group	$\overline{\text{P1}}$ (No. 2)
a, Å	11.0545(19)
b, Å	11.4532(22)
c, Å	20.5023(36)
$\alpha$ , °	86.599(15)
$\beta$ , °	83.524(14)
$\gamma$ , °	62.713(13)
V, Å <sup>3</sup>	2292.11(73)
Z	2
molec wt	856.5
D, g/cm <sup>3</sup>	1.240
$\mu(\text{Mo K}\alpha)$ , mm <sup>-1</sup>	1.204
F(000)	904
2 $\theta$ range, deg	5 - 45
h	0 to +11
k	-10 to +12
$\ell$	-21 to +22
no. of rflns collected	6373
no. of indep rflns	6004 ( $R_{\text{int}} = 1.23\%$ )

**Table 3 (Cont.)**

no. of rflns above $6\sigma$	3711
absorption correction	Semi-empirical
min./max. transmission	0.5747 / 0.7027
weighting scheme, $w^{-1}$	$\sigma^2(F) + 0.0013F^2$
no. of params refined	514
final R indices (all data), %	R = 6.94
	$R_w = 5.65$
R indices ( $6\sigma$ data), %	R = 3.43
	$R_w = 3.62$
goodness of fit	0.93
largest and mean $\Delta/\sigma$	0.002, 0.001
data-to-param ratio	11.7:1
largest diff peak, $e\text{\AA}^{-3}$	0.42
largest diff hole, $e\text{\AA}^{-3}$	-0.42

**Caption to Figures.**

**Figure 1.** Packing diagram for  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$ , viewed down  $a$ . The defined “Molecule 1” (centered about  $0, \frac{1}{2}, 0$ ) lies to the left and “Molecule 2” (centered about  $\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$ ) is in the center of this diagram. (The  $c$ -axis is horizontal and the  $b$ -axis is vertical).

**Figure 2.** Labeling of atoms for “Molecule 1” of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaNHPh}]_2$ . ORTEP2 diagram, 30% probability ellipsoids for non-hydrogen atoms and with all hydrogen atoms artificially reduced.




FIG. 1




Fig. 2.



## TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (1)\*  
Chemistry Division, ONR 331  
800 North Quincy Street  
Arlington, Virginia 22217-5660

Dr. Richard W. Drisko (1)  
Naval Facilities & Engineering  
Service Center  
Code L52  
Port Hueneme, CA 93043

Defense Technical Information Center  
(2)  
8725 John J. Kingman Rd., #0744  
Fort Belvoir, VA 22060-6217

Dr. Eugene C. Fischer (1)  
Code 2840  
Naval Surface Warfare Center  
Carderock Division Detachment  
Annapolis, MD 21402-1198

Dr. James S. Murday (1)  
Chemistry Division, Code 6100  
Naval Research Laboratory  
Washington, D.C. 20375-5320

Dr. John Fischer, Director (1)  
Chemistry Division, C0235  
Naval Air Weapons Center  
Weapons Division  
China Lake, CA 93555-6001

Dr. Peter Seligman (1)  
Naval Command, Control and  
Ocean Surveillance Center  
RDT&E Division  
San Diego, CA 92152-5000

\* Number of copies to forward